

II. *Æquationum Cubicarum & Biquadraticarum, tum Analytica, tum Geometrica & Mechanica, Resolutio Universalis, a J. Colson.*

§. I. *Æquationis Cubicæ* $\left\{ \begin{array}{l} x^3 = 3 p x^2 + 3 q x + 2 r, \\ \text{Universalis} \quad \quad \quad - 3 p^2 + p^3 \\ \quad \quad \quad \quad \quad - 3 p q \end{array} \right.$

Radices Tres sunt,

$$x = p + \sqrt[3]{r + \sqrt{r^2 - q^3}} + \sqrt[3]{r - \sqrt{r^2 - q^3}}$$

$$x = p - \frac{1 - \sqrt{-3}}{2} \times \sqrt[3]{r + \sqrt{r^2 - q^3}} - \frac{1 + \sqrt{-3}}{2} \times \sqrt[3]{r - \sqrt{r^2 - q^3}}$$

$$x = p - \frac{1 + \sqrt{-3}}{2} \times \sqrt[3]{r + \sqrt{r^2 - q^3}} - \frac{1 - \sqrt{-3}}{2} \times \sqrt[3]{r - \sqrt{r^2 - q^3}}$$

Vel ut Calculus Arithmeticus facilius ac paratius evadat, si posueris Binomii irrationalis $r + \sqrt{r^2 - q^3}$ Radicem Cubicam esse $m + \sqrt{n}$, erunt ejusdem *Æquationis* Radices tres $x = p + 2 m$, & $x = p - m + \sqrt{-3 n}$.

Igitur data *Æquatione* quavis Cubica, inter ejus hujusque *Æquationis* Universalis terminos singulos instituenda est comparatio, quo pacto facillime inveniuntur ipsæ p , q , r ; & hisce cognitis, innotescunt *Æquationis* datæ Radices omnes. Hujus vero Solutionis Exempla sint sequentia in Numeris.

I. *Æquationis* Cubicæ $x^3 = 2 x^2 + 3 x + 4$ sit Radix x indaganda. Erit primò juxta præscriptum $3 p = 2$,
14 N five

five $p = \frac{2}{3}$. Secundò $3q - (3p^2) \frac{4}{3} = 3$, five $q = \frac{13}{9}$.

Tertiò $2r (+ \overline{p^2 - 3q \times p}) - \frac{70}{27} = 4$, five $r = \frac{89}{27}$,

& $r^2 - q^3 = \frac{212}{27}$. Et propterea $x = \frac{2}{3} + \sqrt[3]{\frac{89}{27} + \sqrt{\frac{212}{27}}}$

$+ \sqrt[3]{\frac{89}{27} - \sqrt{\frac{212}{27}}}$. Reliquæ duæ Radices sunt impossibiles.

2. In Equatione $x^3 = 12x^2 - 41x + 42$, erit primò $3p = 12$, five $p = 4$. Secundò $3q - (3p^2) 48 = -41$, five $q = \frac{7}{3}$. Tertiò $2r + (\overline{p^2 - 3q \times p}) 36 = 42$,

five $r = 3$; Et inde $r^2 - q^3 = -\frac{100}{27}$. At Binomii furdi

$3 + \sqrt{-\frac{100}{27}} (= r + \sqrt{r^2 - q^3})$ Radix Cubica, per Methodos ex Arithmetica petendas extracta, est $-1 +$

$\sqrt{-\frac{4}{3}} (= m + \sqrt{n},)$ & proinde Radix $x = (p + 2m = 4 - 2 =) 2$, vel etiam $x = (p - m + \sqrt{-3n} = 4 + 1 + (\sqrt{4})^2 =) 7$ vel 3. Vel rursus, ejusdem Binomii

$3 + \sqrt{-\frac{100}{27}}$ Radix alia Cubica (tres enim agnoscit)

est $\frac{3}{2} + \sqrt{-\frac{1}{12}} (= m + \sqrt{n},)$ & proinde Radix

$x = (p + 2m = 4 + 3 =) 7$, & etiam $x = (p - m + \sqrt{-3n} = 4 - \frac{3}{2} + (\sqrt{\frac{1}{4}}) \frac{1}{2} =) 3$ vel 2. Vel denuo,

ejusdem Binomii $3 + \sqrt{-\frac{100}{27}}$ Radix Cubica tertia est

$-\frac{1}{2} - \sqrt{-\frac{25}{12}}, (= m + \sqrt{n},)$ & proinde Radix

$x =$

(2355)

$x = (p + 2m = 4 - 1 =) 3$, atque etiam $x = (p - m \pm \sqrt{m} - 3n = 4 + \frac{1}{2} \pm (\sqrt{\frac{25}{4}}) \frac{5}{2} =) 7$ vel 2.

3. In *Æquatione* $x^3 = -15x^2 - 84x + 100$, erit $p = -5$, $q = -3$, $r = 135$; & Binomii $135 + \sqrt{18252}$ Radix Cubica est $3 + \sqrt{12}$. Igitur Radix $x = -5 + 6 = 1$, & $x = -5 - 3 \pm \sqrt{-36} = -8 \pm \sqrt{-36}$, impossibiles.

4. In *Æquatione* $x^3 = 34x^2 - 310x + 1012$, erit $p = \frac{34}{3}$, $q = \frac{226}{9}$, $r = \frac{5536}{27}$; & Binomii $\frac{5536}{27} + \sqrt{\frac{707560}{27}}$ Radix Cubica est $\frac{16}{3} + \sqrt{\frac{10}{3}}$. Igitur Radix $x = \frac{34}{3} + \frac{32}{3} = 22$, & $x = \frac{34}{3} - \frac{16}{3} \pm \sqrt{-10} = 6 \pm \sqrt{-10}$, impossibiles.

5. In *Æquatione* $x^3 = 28x^2 + 61x - 4048$, erit $p = \frac{28}{3}$, $q = \frac{967}{9}$, $r = -\frac{25010}{27}$; & Binomii $-\frac{25010}{27} + \sqrt{-382347}$. Radix Cubica est $\frac{41}{6} + \sqrt{-\frac{243}{4}}$. Igitur $x = \frac{28}{3} + \frac{41}{3} = 23$, & $x = \frac{28}{3} - \frac{41}{6} \pm (\sqrt{\frac{729}{4}}) \frac{27}{2} = 16$ vel -11 .

6. In *Æquatione* $x^3 = -x^2 + 166x - 660$, erit $p = -\frac{1}{3}$, $q = \frac{499}{9}$, $r = -\frac{9658}{27}$; & Binomii $-\frac{9658}{27} + \sqrt{-\frac{1147205}{27}}$ Radix Cubica est $-\frac{22}{3} + \sqrt{-\frac{5}{3}}$. Igitur $x = -\frac{1}{3} - \frac{44}{3} = -15$, & $x = -\frac{1}{3} + \frac{22}{3} \pm \sqrt{5} = 7 \pm \sqrt{5}$, irrationales.

7. In

(2356)

7. In *Æquatione* $x^3 = 63 x^2 + 99673 x + 9951705$,
erit $p = 21$, $q = \frac{100996}{3}$, $r = 6031680$; & Binomii
 $6031680 + \sqrt[3]{\frac{47887175043136}{27}}$ Radix Cubica est

$183 + \sqrt[3]{\frac{529}{3}}$. Igitur $x = 21 + 366 = 387$, &
 $x = 21 - 183 \pm (\sqrt[3]{529}) 23 = -139$ vel 185 .

Nec fecus in cæteris procedendum: Investigatur autem
Theorema ad modum sequentem. Pono *Æquationis* cu-
jusdam Cubicæ Radicem $z = a + b$, & cubicè multi-
plicando proveniet $z^3 = (a^3 + 3 a^2 b + 3 a b^2 + b^3 =)$
 $a^3 + 3 a b \times a + b + b^3$. Jam loco ipsius $a + b$ valo-
rem ejus z substituendo, fiet $z^3 = 3 a b z + a^3 + b^3$, quæ
est *Æquatio Cubica ex Radice* $z = a + b$ constructa, cui
terminus secundus deest. Ut hæc verò ad formam magis
commodam magisq; concinnam revocenter, sumo *Æqua-*
tionem $z^3 = 3 q z + 2 r$, quæ posthac ipsius $z^3 = 3 a b z$
 $+ a^3 + b^3$ vices gerat. Igitur transmutatione hujus in
illam, fiet primò $3 q = 3 a b$, five $q_3 = a^3 b^3$; & se-
cundò $2 r = a^3 + b^3$, five $2 r a^3 = (a^6 + a^3 b^3 =) a^6 + q^3$.
Et soluta hac *æquatione quadratica*, erit $a^3 = r + \sqrt{r^2 - q^3}$,
& hinc $b^3 = (2 r - a^3 =) r - \sqrt{r^2 - q^3}$: Atque igi-

tur tandem $a = \sqrt[3]{r + \sqrt{r^2 - q^3}}$ & $b = \sqrt[3]{r - \sqrt{r^2 - q^3}}$.
Et propterea in *Æquatione Cubica* $z^3 = 3 q z + 2 r$ erit

Radix $z = (a + b =) \sqrt[3]{r + \sqrt{r^2 - q^3}} + \sqrt[3]{r - \sqrt{r^2 - q^3}}$

At verò hæc Radix reverà triplex est, pro triplici va-
lore quem induere potest & $\sqrt[3]{r + \sqrt{r^2 - q^3}}$ &

$\sqrt[3]{r - \sqrt{r^2 - q^3}}$. Cujusvis enim quantitatis Radix Cu-
bica triplex erit, & ipsius Unitatis Radix Cubica vel
est

est 1, vel $-\frac{1}{2} + \frac{1}{2}\sqrt{-3}$, vel $-\frac{1}{2} - \frac{1}{2}\sqrt{-3}$:

Atque id adeo, propterea quòd harum alicujus Cubus sit

Unitas. Igitur si $1 \times \sqrt[3]{r + \sqrt{r^2 - q^3}}$ aut $\sqrt[3]{r + \sqrt{r^2 - q^3}}$
 $(= \sqrt[3]{1 \times r + \sqrt{r^2 - q^3}} = \sqrt[3]{1} \times \sqrt[3]{r + \sqrt{r^2 - q^3}})$ Ra-
 dicem aliquam [quam supra nominavimus $m + \sqrt{n}$, aut
 $1 \times m + \sqrt{n}$,] Cubi $r + \sqrt{r^2 - q^3}$ designet; ipsæ
 $\frac{-1 + \sqrt{-3}}{2} \times \sqrt[3]{r + \sqrt{r^2 - q^3}}$ & $\frac{-1 - \sqrt{-3}}{2}$
 $\times \sqrt[3]{r + \sqrt{r^2 - q^3}}$ [i. e. $\frac{-1 + \sqrt{-3}}{2} \times m + \sqrt{n}$ &
 $\frac{-1 - \sqrt{-3}}{2} \times m + \sqrt{n}$] alias duas ejusdem Cubi Ra-
 dices designabunt. Similiter & $\sqrt[3]{r - \sqrt{r^2 - q^3}}$,
 $\frac{-1 + \sqrt{-3}}{2} \times \sqrt[3]{r - \sqrt{r^2 - q^3}}$, & $\frac{-1 - \sqrt{-3}}{2}$
 $\times \sqrt[3]{r - \sqrt{r^2 - q^3}}$, [i. e. $m - \sqrt{n}$, $\frac{-1 + \sqrt{-3}}{2}$
 $\times m - \sqrt{n}$, $\frac{-1 - \sqrt{-3}}{2} \times m - \sqrt{n}$] tres Cubicæ Ra-
 dices erunt Apotomes $r - \sqrt{r^2 - q^3}$. Atque has Radices
 debite connecendo, fiet $z = \sqrt[3]{r + \sqrt{r^2 - q^3}}$
 $+ \sqrt[3]{r - \sqrt{r^2 - q^3}}$, [i. e. $z = m + \sqrt{n} + m - \sqrt{n} = 2m$,]
 $z = \frac{-1 + \sqrt{-3}}{2} \times \sqrt[3]{r + \sqrt{r^2 - q^3}} + \frac{-1 - \sqrt{-3}}{2}$
 $\times \sqrt[3]{r - \sqrt{r^2 - q^3}}$, [i. e. $z = \frac{-1 + \sqrt{-3}}{2} \times m + \sqrt{n}$
 $+ \frac{-1 - \sqrt{-3}}{2} \times m - \sqrt{n} = -m + \sqrt{-3}n$,] & $z =$
 $\frac{-1 - \sqrt{-3}}{2} \times \sqrt[3]{r + \sqrt{r^2 - q^3}} + \frac{-1 + \sqrt{-3}}{2} \times \sqrt[3]{r - \sqrt{r^2 - q^3}}$

$$[i. e. z = \frac{-1 - \sqrt{-3}}{2} \times m + \sqrt{n} + \frac{-1 + \sqrt{-3}}{2}$$

$\times m - \sqrt{n} = -m - \sqrt{-3} n]$ quæ tres erunt Radices *Æquationis* Cubicæ $z^3 = 3 q z + 2 r$. Debite autem connectuntur Radices istæ ad modum præcedentem, quippe quæ sic connexæ, & more vulgari in se invicem continueductæ, *Æquationem* $z^3 = 3 q z + 2 r$ restituant. Denique fac $z = x - p$, & *Æquatio* fiet $x^3 - 3 p x^2 + 3 p^2 x - p^3 = 3 q x - 3 p q + 2 r$, quæ universalis est, & cujus Radices evadunt ut supra fuerunt exhibitæ.

Hic obiter notatu dignum est, quod *Æquationis* Cubicæ cujuscunque Radices omnes sint possibiles & reales, quoties Binomii membrum irrationale $\sqrt{r^2 - q^3}$ impossibilitatem in se complectitur; hoc est, quoties q est quantitas affirmativa, & simul cubus ejus major est quadrato ex latere r . At si membrum istud $\sqrt{r^2 - q^3}$ sit possibile, hoc est si q sit quantitas negativa, aut etiam si affirmativæ cubus sit minor quadrato ex latere r , tunc unicam tantum agnoscit *Æquatio* Radicem possibilem & realem, reliquæque duæ erunt impossibiles.

In hoc Theoremate si fiat $p = 0$, hoc est, si desit *Æquationis* terminus secundus, tunc deventum erit ad casum Regularum quæ dicuntur *Cardani*, cujus solutio continetur in præcedentibus.

§. 2. *Æquationis* Biquadraticæ Universalis

$$x^4 = 4 p x^3 + 2 q x^2 + 8 r x + 4s, \\ - 4 p^2 - 4 p q - q^2$$

$$\text{Radices quatuor sunt } x = p - a \pm \sqrt{p^2 + q - a^2} - \frac{2r}{a},$$

$$\& x = p + a \pm \sqrt{p^2 + q - a^2} + \frac{2r}{a}, \text{ Ubi } a^2 \text{ est Radix}$$

$$\text{Æquationis Cubicæ } a^6 = p^2 a^4 - 2 p r a^2 + r^2 \\ + q - s$$

Jam data *Æquatione* quavis Biquadratica, inter ejus hujusque *Æquationis* Universalis terminos singulos instituenda

enda est comparatio, quo pacto citissime inveniuntur ipsæ p, q, r, s; & hisce cognitis, non latebit valor ipsius a, ex Theoremate superiori inveniendus, & tum demum innotescant Æquationis datæ Radices omnes.

Huic Solutioni illustrandæ Exemplum unum aut alterum sufficiat.

1. Æquationis Biquadraticæ $x^4 = 8x^3 + 83x^2 - 162x - 936$ sint Radices extrahendæ. Erit primò juxta præscriptum $4p = 8$, five $p = 2$. Secundò $2q - (4p^2) 16 = 83$, five $q = \frac{99}{2}$. Tertiò $8r - (4pq) 396 = -162$, five $r = \frac{117}{4}$. Quartò $4s - (q^2) \frac{9801}{4} = -936$, five $s = \frac{6057}{16}$. Hinc $p^2 + q = \frac{107}{2}$, $2pr + s = \frac{7929}{16}$, $r^2 = \frac{1368}{19}$, & proinde $a^6 = \frac{107}{2}a^4 - \frac{7929}{16}a^2 + \frac{1368}{16}$. Jam ut Æquatio hæc aliquatenus Cubica in Radices ejus resolvatur, ad Theorema præcedens recurrendum est, in quo erit $p = \frac{107}{2}$, $q = \frac{22009}{144}$, $r = \frac{2903923}{1728}$ & $r^2 - q^3 = -\frac{11940075}{16}$. Atqui Binomii $\frac{2903923}{1728} + \sqrt{-\frac{11940075}{16}}$ Radix Cubica est $-\frac{53}{12} + \sqrt{-\frac{400}{3}}$ & propterea $a^2 = \frac{107}{6} - \frac{53}{6} = 9$, & etiam $a^2 = \frac{107}{6} + \frac{53}{6} \pm (\sqrt{400}) 20 = \frac{169}{4}$ vel $\frac{2}{4}$: Vel quod perinde est, Æquationis præmissæ reverà Cubo-Cubicæ sex Radices sunt $a = \pm 3$, $a = \pm \frac{13}{2}$, & $a = \pm \frac{3}{2}$, quarum quævis indiscriminatim proposito

fito nostro faciet satis. Puta si in præfenti casu fiat $a = 3$, erit juxta Theorema $x = \frac{(p - a + \sqrt{p^2 + q - a^2 - \frac{2r}{a}})}{2} = \frac{2 - 3 \pm \sqrt{4 + \frac{99}{2} - 9 - \frac{39}{2}}}{2} = \frac{-1 \pm (\sqrt{25}) 5}{2} = 4$ vel -6 , & $x = \frac{(p + a + \sqrt{p^2 + q - a^2 + \frac{2r}{a}})}{2} = \frac{2 + 3 \pm \sqrt{4 + \frac{99}{2} - 9 + \frac{39}{2}}}{2} = \frac{5 \pm (\sqrt{64}) 8}{2} = 13$ vel -3 , quæ sunt Æquationis datæ Radices quatuor,

2. In Æquatione $x^4 = 20x^3 + 252x^2 - 6592x + 21312$, erit $p = 5$, $q = 176$, $r = -384$, & $s = 13072$. Hinc $p^2 + q = 201$, $2pr + s = 9232$, & $r^2 = 147456$; & inde $a^6 = 201 a^4 - 9232 a^2 + 147456$.

Jam in Theoremate pro Cubicis erit $p = 67$, $q = \frac{4235}{3}$, & $r = 65219$; eritque Binomii $65219 + \sqrt{\frac{38889307072}{27}}$

Radix Cubica $\frac{77}{2} + \sqrt[3]{\frac{847}{12}}$. Igitur $a^2 = 67 + 77 = 144$, five $a = 12$; & proinde $x = \frac{5 - 12 \pm \sqrt{25 + 176 - 144 + 64}}{2} = \frac{-7 \pm (\sqrt{121}) 11}{2} = 4$ vel -18 , & $x = \frac{5 + 12 \pm \sqrt{25 + 176 - 144 - 64}}{2} = \frac{17 \pm \sqrt{-7}}{2}$, impossibiles.

Hujus autem Theorema's Inventio est hujusmodi, Ex duarum Æquationum Quadraticarum $z^2 + 2az - b = 0$, & $z^2 - 2az - c = 0$ in se invicem multiplicatione, Æquationem conficio Biquadraticam $z^4 = 4a^2 + b + c \times z^2 + 2ac - 2ab \times z - bc$, cui terminus secundus deest, quamque huc Æquationi $z^4 = ez^2 + fz + g$ statuo æquipollere. Unde primo $4a^2 + b + c = e$ five $b = e - 4a^2 - c$. Secundò $2ac - 2ab = f$, hoc est, $2ac - 2ae + 2a^3 + 2ac = f$, five $c = \frac{f}{4a} + \frac{e}{2} - 2a^2$,

& inde $b = (e - 4a^2 - c) - \frac{f}{4a} + \frac{e}{2} - 2a^2$. Ter-

tiò $-bc = g$, five $-\frac{f^2}{16a^2} + \frac{e^2}{4} - 2ea^2 + 4a^4 = -g$,

hoc est, $a^4 = \frac{1}{2} ea^2 - \frac{1}{4} ga^2 - \frac{1}{16} ea^2 + \frac{f^2}{64}$, quæ

Æquatio quasi Cubica est, ex Radice a^2 & notis vel as-
sumptis e, f, g conflata. Ea verò Radix per Theorema
superius exhiberi potest, & eodem Calculo innotescant
ipsæ b & c . At Æquationum $z^3 + 2az - b = 0$ &
 $z^3 - 2az - c = 0$ Radices sunt $z = -a \pm \sqrt{a^2 + b}$

& $z = a \pm \sqrt{a^2 + c}$, five $z = -a \pm \sqrt{\frac{1}{2}e - a^2 - 4a^2}$,

& $z = a \pm \sqrt{\frac{1}{2}e - a^2 + \frac{f}{4a}}$, quæ proinde erunt Radices
Æquationis $z^4 = ez^2 + fz + g$; cognita videlicet a vel a^2
ex Æquatione $a^6 = \frac{1}{2}ea^4 - \frac{1}{4}ga^2 - \frac{1}{16}ea^2 + \frac{f^2}{64}$. Jam ut

Æquatio ista evadat universalis, & omnibus suis terminis
instructa, fac. $z = x - p$, eritque $x^4 - 4px^3 + 6p^2x^2$
 $- 4p^3x + p^4 = ex^2 - 2pex + p^2e + fx - fp + g$,

item & $x = p - a \pm \sqrt{\frac{1}{2}e - a^2 - \frac{f}{4a}}$, & $x = p + a \pm$
 $\sqrt{\frac{1}{2}e - a^2 + \frac{f}{4a}}$. Tandem concinnitatis & compendii

gratiâ, fac. $e = 2q + 2p^2$ & $f = 8r$; tum $x^4 - 4px^3$
 $+ 4p^2x^2 = 2qx^2 - 4pqx + 2p^2q + p^4 + 8rx - 8pr + g$,

$x = p - a \pm \sqrt{p^2 + q - a^2 - \frac{2r}{a}}$, $x = p + a \pm$
 $\sqrt{p^2 + q - a^2 + \frac{2r}{a}}$, & $a^6 = p^2 + q + a^4 - \frac{1}{4}g + \frac{1}{4}p^4$

$+ \frac{1}{2}p^2q - \frac{1}{4}q^2 + a^2 + r^2$. Denique fac $g = 4s - q^2$
 $+ 8pr - p^4 - 2p^2q$, & fiunt Æquationes præcedentes
 $x^4 = 4px^3 + 2qx^2 + 8rx + 4s$ & $a^6 = p^2a^4 - 2pra^2 + r^2$
 $- 4p^2 - 4pq - q^2 + q - s$

Scilicet omnia evadunt ut supra sunt posita.

§ 3. Hactenus de *Æquationum Cubicarum & Biquadraticarum Resolutione Analytica*. Quoniam autem earundem *Effectio Geometrica* per Parabolam vulgò tradi solet, & nonnullis in pretio est, ipsam *construere*, & quidem universalius, non pigebit hic exhibere.


Data *Æquatione* quavis vel Cubica vel Biquadratica, instituenda est comparatio inter terminos ejus, terminosque respondentes hujus *Æquationis*

$$x^4 = \frac{2p}{q} x^3 + \frac{4pr}{q} x^2 + \frac{2p^2}{q} x + p^2, \text{ quo pacto facile satis}$$

— 4r	— 4r ²	— $\frac{2p^2}{q}$	— q ²
	+ 2s	+ 4rs	— s ²
	— 1	— 2q	+ t ²

eruentur ipsæ p, q, r, s, t; earum interim unâ aliquâ ut-
cunque pro lubitu assumptâ. Tum in Parabola quavis
data AVB, cujus Vertex principalis V, Axis VS, & Axi

(2363)



perpendicularis VT; capiatur VS = p versus interiora Parabolæ, & in angulo SVT inscribatur ST = q, quæ producta Parabolam secet in punctis binis N & O. Bifecetur ON in M, & per M agatur MA Axi parallela & Parabolæ occurrens in A. Ipsi ON parallela ducatur AL, ut sit AL Latus rectum Parabolæ ad Diametrum AM, sitque hæc eadem Unitas. In AL (utrinque si opus est producta) capiatur AG = r, & à puncto G ducatur GR Axi parallela, quæ Parabolam secet in B, à quo capiatur BR = s. A novissime invento puncto R ducatur RE ipsi VT parallela & æqualis, quæ sinistram versus jaceat respectu ipsius R si q sit quantitas affirmativa, at versus dextram si q sit negativa. Atque idem de ipsis AG & BR intelligatur, quæ ad contrarias itidem partes duci debent, si modò valores ipsarum r & s prodeant negativi. Denique Centro E & Radio EC = t describatur Circulus CK^æ, qui Parabolam in totidem secabit punctis, quot sunt Æquationis datæ Radices reales. Etenim à punctis istis C, K, &c. ducantur CP, kn, &c. ipsi ST parallela, & ad rectam GR (si opus est productam) terminatæ, eritque harum quævis x, seu Æquationis datæ Radix quæsitæ; eæ scilicet ad dextram jacentes erunt Radices affirmativæ, quæ verò ad sinistram sunt positæ erunt Radices negativæ. Punctum contactûs, siquod fuerit, hic sumitur pro intersectionis punctis duobus ad invicem vicinissimis.

Inter Æquationes Cubicas & Biquadraticas ita constructas hoc tantum intercedit discriminis, quod in prioribus, ob terminum ultimum in præcedente Æquatione deficientem, semper sit $p^2 - q^2 - s^2 + t^2 = 0$, sive $t = \sqrt{s^2 + q^2 - p^2}$. Igitur Centro E & Radio EB ($= \sqrt{BRq + (ERq)STq - VSq}$) descripto Circulo CK^æ, Radicum una CP in priori constructione in nihilum abit.

Hæc autem demonstrantur ad modum sequentem. Momentibus jam constructis, & producta CP si opus est, donec secat AM in H, erit CH Ordinata Parabolæ ad Diametrum

metrum AH, & proinde $CHq = AL \times AH = AH$, ob
 $AL = 1$. At $CH = CP + AG$, & $AH = GB + BP$, &
 propterea $CPq + 2AG \times CP + AGq = GB + BP$; sed
 ob naturam Parabolæ erit $AGq = GB$, unde $CPq + 2AG$
 $\times CP = BP$. Jam à puncto C ad ipsam BP demittatur
 norma s CD, quæ occurrat etiam ipsi EI, ad BP actæ pa-
 rallelæ, in puncto I. Propter similia Triangula CDP &

$$TVS, \text{ erit } DP = \frac{VS \times CP}{ST} \text{ \& } CD = \frac{VT \times CP}{ST}, \text{ \& pro-}$$

$$\text{inde } CPq + 2AG \times CP = BP = DP + BD = \frac{VS \times CP}{ST}$$

$$+ BR - IE, \text{ five } CPq + 2AG \times CP - \frac{VS}{ST} CP - BR$$

$$= -IE. \text{ At } IEq = CEq - CIq = CEq - CDq$$

$$- VTq - 2CD \times VT = CEq - \frac{VTq \times CPq}{STq} - VTq$$

$$- \frac{2VTq \times CP}{ST} = (\text{ob } VTq = STq - SVq) CEq - CPq$$

$$+ \frac{SVq}{STq} CPq - STq + SVq - 2ST \times CP + \frac{2SVq}{ST} CP,$$

quæ igitur æqualis erit Quadrato ex Latere $CPq + 2AG$
 $\times CP - \frac{VS}{ST} CP - BR$. Atque hæc Æquatio ad termi-

nos p, q, r, s, t revocata ipsissima sit Æquatio proposita.

Hinc liquet, quòd eadem quævis Æquatio Biquadratica
 innumeras per Parabolam constructiones fortiri possit, pro
 indefinito valore quantitatis istius, quam ad arbitrium assu-
 mi posse jam diximus. Sed casus est simplicissimus faciendo
 $VS = p = 0$, & migrat constructio, si rem ipsam spectes,
 in vulgarem istam, in qua Radicum repræsentatrices
 rectæ CP, &c. sunt ad Axem perpendiculares. Æquatio
 autem fit $x^4 = -4rx^3 - 4r^2x^2 + 4rsx - q^2$, quæ facile

$$\begin{array}{rcccc} & + 2s & = & 2q & - s^2 \\ & - 1 & & & + 1^2 \end{array}$$

construitur ut supra.

§ 4. Sed ne Parabolæ descriptio Organica difficilis nimium videatur, in promptu est Artificium quoddam Mechanicum, op^e Fili penduli pondere instructi peractum, cujus auxilio quam exactissime & facillime Æquatio novissima construi potest, & proinde Æquationum quarumcunque Cubicarum & Biquadraticarum Radices inveniri; idque sine ullo linearum ductu nisi Rectarum & Circuli. Constructio autem, quam appellare libet *Mechanicam*, est ad hunc modum.

Contra Parietem erectum, vel planum aliud quodvis Horizonti perpendiculare, ad punctum aliquod *F* suspendatur filum tenuissimum & flexile *FP*; pondere quovis *P* ad extremitatem *P* appenso. In hoc filo notetur punctum aliquod *N*, à puncto suspensionis *F* satis remotum; vel filo parvulus, si id mavis, innectatur Nodus *N*. Et sumpta utcunque *NO* pro Unitate, ad punctum medium *A* ducatur (in plano prædicto) recta *AQ* Horizonti parallela, & utrinque quantum satis producta. Hisce generaliter paratis, pro particulari jam applicatione fac *AQ* = *r*, ipsis *q*, *r*, *s*, *t*, ut sæpius inculcatum, vel Arithmeticè vel Geometricè, pro datæ cujusvis Æquationis

ex-

quæ erunt *Æquationis* datæ Radices omnes reales ; hæ nempe ad dextram erunt Radices affirmativæ, illæ verò ad sinistram Radices negativæ. Demonstratio est manifesta ex præcedentibus, habita tantùm ratione Parabolæ per puncta B, C, c, x, x tranſeantis. Nam poſito F foco Parabolæ, (cujus diſtantiâ à Vertice aſt $\frac{1}{2}$ ON,) notum eſt quod lineæ omnes ut FB + BQ, FC + CD, &c, eandem ubique conſiciant ſummam.

Atque ex principiis hic poſitis proclive erit Inſtrumentum haud inconcinnum & quantumvis accuratum fabricari, cujus beneficio hujusmodi *Æquationum* quarumcunque Radices nullo fere negotio inveniri poſſint, & præ oculis exhiberi. Hoc autem quilibet, ſi id Curæ ſit, variis modis pro ingenio ſuo efficere poteſt, & de his jam ſatis.

III. *Æquationum* quarundam Potestatis tertiæ, quintæ, ſeptimæ, nonæ, & ſuperiorum, ad infinitum uſque pergendo, in terminis finitis, ad inſtar Regularum pro Cubicis quæ vocantur Cardani, Reſolutio Analytica.

Per Ab. De Moivre, R. S. S.

Si n Numerus quicunque, y quantitas incognita, ſive *Æquationis* Radix quæſita, ſitque a quantitas quævis omnino cognita, ſive ut vocant Homogeneum Comparationis : Atque horum inter ſe relatio exprimatur per *Æquationem*

$$ny + \frac{nn - 1}{2 \times 3} ny^3 + \frac{nn - 1}{2 \times 3} \times \frac{nn - 9}{4 \times 5} ny^5 + \frac{nn - 1}{2 \times 3} \times \frac{nn - 9}{4 \times 5} \times \frac{nn - 25}{6 \times 7} ny^7, \&c. = a$$